



JYOTHISHMATHI INSTITUTE OF TECHNOLOGY & SCIENCE

Sub: HEAT TRANSFER

III B TECH-II SEM (A.Y:2018-19)

RADIATION HEAT TRANSFER

BY

P.DEVENDAR (Asst.Prof)

Department Of Mechanical Engineering.

Blackbody Radiation

- ▶ Blackbody – a perfect emitter & absorber of radiation
- ▶ Emits radiation uniformly in all directions – no directional distribution – it's diffuse
- ▶ Joseph Stefan (1879)– total radiation emission per unit time & area over all wavelengths and in all directions:

$$E_b = \sigma T^4 \left(\text{W} / \text{m}^2 \right)$$

- ▶ σ =Stefan–Boltzmann constant = 5.67×10^{-8}
 $\text{W} / \text{m}^2 \text{K}^4$

Planck's Distribution Law

- ▶ Sometimes we're interested in radiation at a certain wavelength
- ▶ Spectral blackbody emissive power ($E_{b\lambda}$) = “amount of radiation energy emitted by a blackbody at an absolute temperature T per unit time, per unit surface area, and per unit wavelength about the wavelength λ .”

Planck's Distribution Law

- ▶ For a surface in a vacuum or gas

$$E_{b\lambda}(T) = \frac{C_1}{\lambda^5 [\exp(C_2/\lambda T) - 1]} \quad (\text{W} / \text{m}^2 \cdot \mu\text{m})$$

where


$$C_1 = 2\pi hc_o^2 = 3.742 \times 10^8 \text{ W} \cdot \mu\text{m}^4 / \text{m}^2$$

$$C_2 = hc_o / k = 1.439 \times 10^4 \mu\text{m} \cdot \text{K}$$

$$k = 1.3805 \times 10^{-23} \text{ J/K} = \text{Boltzmann's constant}$$

- ▶ Other media: replace C_1 with C_1/n^2
- ▶ Integrating this function over all λ gives us the equation for E_b .

Radiation Distribution



- ▶ Radiation is a continuous function of wavelength
- ▶ Radiation increases with temp.
- ▶ At higher temps, more radiation is at shorter wavelengths.
- ▶ Solar radiation peak is in the visible range.

Wien's Displacement Law

- ▶ Peak can be found for different temps using Wien's Displacement Law:

$$(\lambda T)_{\max \text{ power}} = 2897.5 \mu\text{m} \cdot \text{K}$$

- ▶ Note that color is a function of absorption & reflection, not emission


Blackbody Radiation Function

- ▶ Use blackbody radiation function, F_λ

$$F_\lambda(T) = \frac{\int_0^\lambda E_{b\lambda}(T) d\lambda}{\sigma T^4}$$

- ▶ If we want radiation between λ_1 & λ_2 ,

$$F_{\lambda_1 - \lambda_2}(T) = F_{\lambda_2}(T) - F_{\lambda_1}(T)$$



Surface Emission

Spectral, directional emissivity

$$\varepsilon_{\lambda,\theta}(\lambda, \theta, \phi, T) \equiv \frac{I_{\lambda,e}(\lambda, \theta, \phi, T)}{I_{\lambda,b}(\lambda, T)}$$

Total, directional emissivity

$$\varepsilon_{\theta}(\theta, \phi, T) \equiv \frac{I_e(\theta, \phi, T)}{I_b(T)}$$

Spectral, hemispherical emissivity

$$\varepsilon_{\lambda}(\lambda, T) \equiv \frac{E_{\lambda}(\lambda, T)}{E_{\lambda,b}(\lambda, T)}$$

$$\varepsilon_{\lambda}(\lambda, T) = \frac{\int_0^{2\pi} \int_0^{\pi/2} I_{\lambda,e}(\lambda, \theta, \phi, T) \cos \theta \sin \theta d\theta d\phi}{\int_0^{2\pi} \int_0^{\pi/2} I_{\lambda,b}(\lambda, T) \cos \theta \sin \theta d\theta d\phi}$$

Surface Emission

Spectral, hemispherical emissivity

$$\varepsilon_{\lambda}(\lambda, T) \equiv \frac{E_{\lambda}(\lambda, T)}{E_{\lambda,b}(\lambda, T)}$$

Substituting spectral emissive power

$$\varepsilon_{\lambda}(\lambda, T) = \frac{\int_0^{2\pi} \int_0^{\pi/2} I_{\lambda,e}(\lambda, \theta, \phi, T) \cos \theta \sin \theta d\theta d\phi}{\int_0^{2\pi} \int_0^{\pi/2} I_{\lambda,b}(\lambda, T) \cos \theta \sin \theta d\theta d\phi}$$



Surface Emission

Total, hemispherical emissivity


$$\varepsilon(T) \equiv \frac{E(T)}{E_b(T)}$$

$$\varepsilon(T) = \frac{\int_0^\infty \varepsilon_\lambda(\lambda, T) E_{\lambda,b}(\lambda, T) d\lambda}{E_b(T)}$$

Total, directional emissivity
Normal emissivity predictable




Total, normal emissivity



Temperature dependence of the total, normal emissivity ϵ_n of selected

Absorption, Reflection, and Transmission

$$G_{\lambda} = G_{\lambda,\text{ref}} + G_{\lambda,\text{abs}} + G_{\lambda,\text{tr}}$$



Absorption, reflection, and transmission processes associated with a semitransparent medium.

Absorptivity

Spectral, directional absorptivity

$$\alpha_{\lambda,\theta}(\lambda, \theta, \phi) \equiv \frac{I_{\lambda,i,\text{abs}}(\lambda, \theta, \phi)}{I_{\lambda,i}(\lambda, \theta, \phi)}$$

Spectral, hemispherical absorptivity

$$\alpha_{\lambda}(\lambda) \equiv \frac{G_{\lambda,\text{abs}}(\lambda)}{G_{\lambda}(\lambda)}$$

Total, hemispherical absorptivity

$$\alpha \equiv \frac{G_{\text{abs}}}{G}$$

Reflectivity

Spectral, directional reflectivity

$$\rho_{\lambda,\theta}(\lambda, \theta, \phi) \equiv \frac{I_{\lambda,i,\text{ref}}(\lambda, \theta, \phi)}{I_{\lambda,i}(\lambda, \theta, \phi)}$$



Spectral, hemispherical reflectivity

$$\rho_{\lambda}(\lambda) \equiv \frac{G_{\lambda,\text{ref}}(\lambda)}{G_{\lambda}(\lambda)}$$

Total, hemispherical reflectivity

$$\rho \equiv \frac{G_{\text{ref}}}{G}$$

Reflectivity



Diffuse and specular reflection.

Transmissivity

Spectral, hemispherical transmissivity

$$\tau_{\lambda} = \frac{G_{\lambda, \text{tr}}(\lambda)}{G_{\lambda}(\lambda)}$$

Total, hemispherical reflectivity


$$\tau = \frac{G_{\text{tr}}}{G}$$

Special Considerations

Transparent medium


$$\rho + \alpha + \tau = 1$$

Opaque

$$\rho + \alpha = 1$$


Kirchhoff's Law

$$\varepsilon = \alpha$$



Radiative exchange in an isothermal enclosure.

$$\frac{E_1(T_s)}{\alpha_1} = E_b(T_s)$$

No real surface can have an emissive power exceeding that of a black surface at the same temperature.

ANY QUIRIES ???